

Update on ESTIM-rHTN in France

Pr Patrick Rossignol

Inserm 1433

Clinical Investigation Center

Hypertension and Heart Failure Unit

CHRU Nancy & University of Lorraine, Nancy, France

Association Lorraine pour le Traitement de l'Insuffisance Rénale (ALTIR)

BAROSTIM THERAPY SUMMIT

September 30th, 2017 • Radisson Blu, Berlin, Germany



BAROSTIM
THERAPY™

Agenda

- Background
- Trial Design
- Trial Status




BAROSTIM
THERAPY™

Background

- Rheos double blind pivotal study (Bisognano et al, JACC 2011), preliminary evidence of TOD (LVM) protection (Bisognano, JACC 2011), long-term BP effects (Bakris, JASH, 2012)
- Preliminary evidence of efficacy/safety of the second generation BAROSTIM NEO™ device (Hoppe, JASH 2012)
- **Possible BP-pressure lowering effect of the implantation per se (Bakris, JASH 2012)**

Phase-3 Long-Term Data in Resistant HTN



IS A « COMPLETE »
SHAM APPROPRIATE
FOR FUTURE
STUDIES NOW?

Bakris et al., J Am Soc Hypertens 2012, 6(2):152-158

Background

- Rheos double blind pivotal study, preliminary evidence of TOD (LVM) protection, long-term BP effects
- Preliminary evidence of efficacy/safety of the second generation BAROSTIM NEO™ device
- Possible BP-pressure lowering effect of the implantation per se
- **Need for a new RCT with the BAROSTIM NEO™ device, to enable a health economics assessment in the French healthcare system, prior to reimbursement**

Agenda

- Background
- Trial Design
- Trial Status



BAROSTIM
THERAPY™

ESTIM-rHTN has been granted by the French Ministry of Health & is sponsored by Nancy University Hospital, France

ESTIM-rHTN:

medicoEconomic evaluation of baroreceptor STIMulation
for the treatment of resistant HyperTensioN

A multicenter, randomized, PROBE design trial of baroreceptor stimulation with BAROSTIM NEO™ vs. usual care

ClinicalTrials.gov Identifier:
NCT02364310

P. Rossignol, PI
M. Azizi, co-PI

Main Inclusion Criteria

- 18-85 year-old men or women
- Resistant hypertension on 4 medications including thiazide or thiazide-like at appropriate doses + spironolactone (unless spiro intolerance)
- Essential hypertension
- eGFR \geq 30 mL/min/1,73 m²
- No carotid condition associated with a contra-indication to use BAROSTIM NEO™

Primary Objective

- Medico-economic assessment:
cost-effectiveness

Main Evaluation Criteria

Medico-economic assessment:


Mechanism	Criteria	Unit
Clinical Trial	ICER	Cost / Diurnal SBP mm Hg on ABPM after 12 months
Modeling	Incremental ratio	Cost / Living year saved

Secondary Outcome Criteria

Comparison
with usual
care:

Criteria	6 months	12 months
SBP, DBP, mean BP 24 h, diurnal, nocturnal on ABPM	X	X
Casual BP, PWV, PPc, LVM, LV function on echocardiography	X	X
Evolution of eGFR (MDRD) and of microalbuminuria	X	X
Antihypertensive regimen	X	X
EuroQol5D	X	X
Global cardiovascular risk	X	X
Compliance		
Cardiovascular, cerebrovascular events		
Tolerance : morbidity at 1 month post surgery		

Experimental Design: Probe



Trial Organisation

Steering Committee:

- **Chair:**
- **Co-chair:**
- **Methodologist, statistician:**
- **Medico-economic assessment:**
- **Vascular surgeon:**
- **Cardiologist, HF specialist:**
- **Advisor:**

Pr Patrick Rossignol, Nancy
Pr Michel Azizi, Paris
Pr Gilles Chatellier, Paris
Dr Karine Chevreul (URC-Eco, Paris)
Dr Jean-Marc Alsac, Paris
Pr Faiez Zannad, Nancy
Pr George Bakris, Chicago, USA

DSMB:

- **Pr Pascal Desgranges:**
- **Pr Pascal Bousquet:**
- **Dr Catherine Cornu:**

Vascular surgeon, Hôpital Henry Mondor, Créteil
Cardiologist, Pharmacologist, Université de Strasbourg
Methodologist, CIC, CHU de Lyon



www.inicrct.org

The screenshot shows the INI-CRCT website. At the top, there is a red header bar with the INI-CRCT logo on the left and the Fcrin logo on the right. Below the header, the main navigation menu includes links for HOME, OUR NETWORK, OUR RESEARCH, PARTICIPATE IN OUR RESEARCH PROGRAMS, FELLOWS, and SUPPORT US. The main content area features a large image of a brain with blue and green patterns. Below the image, there is a search bar and a "More information" button. On the right side of the page, there is a "News" section with a blue header and some text.

Agenda


- Background
- Trial Design
- Trial Status



BAROSTIM
THERAPY™

All 16 study sites are activated and can randomise patients





The double challenge of resistant hypertension and chronic kidney disease

Patrick Rossignol, Ziad A Massy, Michel Azizi, George Bakris, Eberhard Ritz, Adrian Covic, David Goldsmith, Gunnar H Heine, Kitty JJager, Mehmet Kanbay, Francesca Mallamaci, Alberto Ortiz, Raymond Vanholder, Andrzej Wiecek, Carmine Zoccali, Gérard Michel London, Bénédicte Stengel, Denis Fouque, for the ERA-EDTA EURECA-m working group, Red de Investigación Renal (REDINREN) network, and Cardiovascular and Renal Clinical Trialists (F-CRIN INI-CRCT) network



BAROSTIM
THERAPY™

Thank you.



BAROSTIM
THERAPY™